

EGU2020-20531

<https://doi.org/10.5194/egusphere-egu2020-20531>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



Estimating the Above Ground Biomass of Brazilian Savanna using multi-sensor approach

Polyanna da Conceição Bispo^{1,2}, Pedro Rodriguez-Veiga^{2,3}, Barbara Zimbres⁴, Sabrina do Couto de Miranda⁵, Cassio Henrique Giusti Cezare⁶, Sam Fleming⁷, Francesca Baldacchino⁷, Julia Zanin Shimbo⁴, Ane Auxiliadora Costa Alencar⁴, Iris Roitman⁸, Mercedes Bustamante⁸, Ana Maria Pacheco-Pascagaza², Yaqing Gou², John Roberts², Valentin Louis², Kirsten Barret², Iain Woodhouse^{7,9}, Eráclito Sousa-Neto¹⁰, Jean P.H.B. Ometto¹⁰, and Heiko Balzter^{2,3}

¹Department of Geography, School of Environment, Education and Development, University of Manchester, Manchester, United Kingdom (polyanna.bispo@manchester.ac.uk)

²Centre for Landscape and Climate Research, School of Geography Geology and the Environment, Department of Geography, University of Leicester, Leicester, United Kingdom

³National Centre for Earth Observation at the University of Leicester, Leicester, United Kingdom

⁴Amazon Environmental Research Institute (IPAM), Brasília, Brazil

⁵State University of Goiás (UEG), Palmeiras de Goiás, Brazil

⁶Federal University of Goiás (UFG), Goiânia, Brazil

⁷Carbomap Ltd., Edinburgh, United Kingdom

⁸Department of Ecology, Universidade de Brasília (UNB), Brazil

⁹School of Geosciences, University of Edinburgh, Edinburgh, United Kingdom

¹⁰Earth System Science Center of the National Institute for Space Studies (CCST-INPE), São José dos Campos, Brazil

The Brazilian Savanna, known as Cerrado (Cerrado sensu lato (s.l.)), is the second largest biome in South America. It comprises different physiognomies due to variations of soil, topography and human impacts. The gradients of tree density, tree height, above ground biomass (AGB) and wood species cover vary according to the Cerrado formation, ranging from different grassland formations (Campo limpo, campo sujo), savanna intermediary formations (Campo cerrado and Cerrado sensu stricto - s.s) and forest formations (Cerradão, Mata ciliar, Mata de galeria and Mata Seca).

Although the carbon stock in Cerrado is lower than in the Brazilian Amazon, the conversion of this biome to other types of land use is occurring much faster. In the last ten years, the degradation of Cerrado forest was the second largest source of carbon emissions in Brazil. Therefore, effective methods for assessing and monitoring aboveground woody biomass and carbon stocks are needed. A multi-sensor Earth observation approach and machine learning techniques have shown potential for the large-scale characterization of Cerrado forest structure. The aim of this study is to present a method to estimate the AGB of an area of the Brazilian Cerrado using ALOS-PALSAR (L-band SAR), Landsat, LIDAR (Light Detection And Ranging) and field datasets. Field data consisted of 15 plots of 1 ha area located in Rio Vermelho watershed in Goiás-State (Brazil). We used a 2-step AGB estimation (i) from the field AGB using LIDAR metrics and (ii) from LIDAR-AGB to satellite Earth

Observation scales following a Random Forest regression algorithm. The methodology to estimate ABG of Cerrado Stricto Sensu vegetation is part of the Forests 2020 project which is the largest investment by the UK Space Agency, as part of the International Partnerships Programme (IPP), to support in the improvement of the forest monitoring in six partner countries through advanced uses of satellite data.