



MULTI-ANNUAL CHANGES OF GLACIAL LAKES IN PERU AND THEIR LINKS TO CLIMATE CHANGE AND GLOFS

Turpo Cayo E.(4,5); Jara Infantes, H.(1); Melgarejo, E.(1); Tinoco Meyhuay T.(3); Torres, J (1); Caballero, A.(1); Villafane, H.(1); Araujo,J.(1); Yarleque, C.(1); Garay, D. (1); Stephan,H (2); Wood,J (2); Glasser, N.(2); Ryan W.(2); Bennett, J.(2); Shannon, S.(2); Espinoza, R.(5) and De Almeida, C.(6,5)

(1) National Research Institute for Glaciers and Mountain's Ecosystems (INAIGEM); (2) University of Exeter; (3) Santiago Antúnez de Mayolo National University (UNASAM); (4) Instituto de Geografía y Geología (IGG); (5) Universidad Nacional Mayor de San Marcos (UNMSM); (6) National Institute of Water Research (INIA)

ABSTRACT


The effects of climate change are causing a profound reduction in glacier's coverage in the Cordilleras of the Peruvian Andes, giving rise to the development and growth of proglacial lakes. These glacial lakes can represent a significant hazard in the form of Glacier Lake Outburst Floods (GLOF) that have the potential to impact vulnerable populations and infrastructure located downstream of glaciated areas. In this study, we present a long-term spatio-temporal analysis of glacial lakes across the Peruvian Andes and address possible connections with GLOF events and climate change. Accessed and processed using Google Earth Engine and QGIS, this analysis was conducted using Landsat 5, 7 and 8 satellite data (Surface Reflectance Tier 1) acquired between 1984 and 2019. We detected and estimated glacial lake surface area changes using a robust semi-automated

METHODOLOGY

The pre-processing and image processing procedures for the multi-temporal mapping of glacial lakes were based on an adaptation of the Mapbiomas project methodology (MapBiomas 2017) and other references (Pekel et al. 2016, Cayo Turpo et al. 2019, Souza et al. 2019). The main characteristics of the proposed methodology include:

- usage of the Landsat collection (30 m spatial resolution),
- pixel-based image processing (empirical decision tree),
- organized by Cordilleras,
- cloud-based computing (Google Earth Engine),
- multi-year validation and field survey.

Figure 1 Methodological flowchart



RESULTS

According to Figure 4, we showed the evolution of the surface extension of the glacial lakes in the 20 Cordilleras of the Peruvian Andes in the period 1984-2019; where the La Viuda Cordillera has the greatest surface extension. The Cordilleras that presented the greatest increase in the surface of glacial lakes are Ampato, Volcánica, Apolobamba, and Chonta, while the Huytapallana and Huagorucho Cordilleras have a decreasing trend. In the case of the Barroso Cordillera, a significant increase was found, related to the activity of damming and regulation of lakes. In the rest of the Cordilleras, there are less representative changes, however, in the lakes individual analysis, important changes were found in most of the studied Cordilleras of the Peruvian Andes.

Figure 3. Classification of glacial lakes based on empirical decision tree.





Figure 4. Glacial lakes area evolution in the 20 Cordilleras of the Peruvian Andes.



DISCUSSION

There are several methodologies to classify water bodies in satellites images, of which empirical decision trees are one most used due to their easiness of use and implementation (Souza et al. 2019, MapBiomas Amazonia 2019). However, empirical decision trees can be subjective in the parameters setting at each decision node, leading to temporally inconsistent results, requiring the application of different post-classification filters. For this research, it was decided to use empirical decision trees due to their superior quality when compared to other classifiers such as Random Forest in the Andes mountain range, where there is a general confusion between the classification of water and relief shadows.

The results of the glacial lakes classification in the period from 1984 to 2019 have overall accuracy greater than 85%. This indicates that the methodology is appropriate for the study area. It ought to be noted that the quality of the classification depends on the variables than are used

CONCLUSIONS

This study applied a methodology based on empirical decision trees, to the entire Andes mountains. The classification was based on Landsat satellite images and was carried out in the 20 Peruvian Cordilleras in the period 1984-2019. In general, in all Andes mountains, the overall accuracy surpassed 85%. This indicates that the results of this study are an important source of information for the analysis of glacial lakes, their evolution, and trends in the latest three decades.

MULTI-ANNUAL CHANGES OF GLACIAL LAKES IN PERU AND THEIR LINKS TO CLIMATE CHANGE AND GLOFS

MULTI-ANNUAL CHANGES OF GLACIAL LAKES IN PERU AND THEIR LINKS TO CLIMATE CHANGE AND GLOFS
 Turpo Cayo E.(4,5); Jara Infantes, H.(1); Melgarejo, E.(1); Tinoco Meyhuay T.(3); Torres, J (1); Caballero, A.(1); Villafane, H.(1); Araujo,J.(1); Yarleque, C.(1); Garay, D. (1); Stephan,H (2); Wood,J (2); Glasser, N.(2); Ryan W.(2); Bennett, J.(2); Shannon, S (2); Espinoza, R.(5) and De Almeida, C.(6,5)

(1) National Research Institute for Glaciers and Mountain's Ecosystems (INAIGEM); (2) University of Exeter; (3) Santiago Antúnez de Mayolo National University (UNASAM); (4) Institute of the Common Good (IBC); (5) La Molina National Agrarian University, Doctoral Program in Water Resources (PDRH); (6) National Institute for Space Research (INPE)

ABSTRACT
 The effects of climate change are causing a profound reduction in glacier coverage in the Cordillera of the Peruvian Andes, giving rise to the development and growth of glacial lakes. These glacial lakes can represent a significant hazard for the town of Glacier Lake (Cajamarca) due to the generation of rapid meltwater pulses and subsequent limited development of glacial lakes. In this regard, we present a long-term spatial-temporal evolution of glacial lakes across the Peruvian Andes and address possible connections with GLOFS events and climate change. We used an open-access satellite-based Landsat 5 image and GTOPO30, and we use a combined using Landsat 5, 30-m GTOPO30

RESULTS
 According to Figure 1, we observed the evolution of the spatial extension of the glacial lakes in the 20 Cordilleras of the Peruvian Andes in the period 1984-2019, using the Landsat 5 satellite-based data. The Cordillera that presented the greatest increase in the surface of glacial lakes are 'Cajamarca', 'Vilcabamba', 'Apurimac', and 'Cuzco', while the 'Huancabamba' and 'Huancabamba' Cordillera areas decreasing in total. In the case of the Huancabamba, a significant increase was found, related to the activity of damming and regulation of lakes. In the rest of the Cordillera, there are less pronounced changes. However, in the inter-annual analysis, important changes were found in most of the studied Cordilleras of the Peruvian Andes.

Figure 1. Classification of glacial lakes based on temporal extension

DISCUSSION
 There are several methodologies to classify water bodies in satellite images, of which temporal detection was one of the most effective in this context and was applied to the Landsat 5 images of the 20 CD. Additionally, however, the temporal analysis of the data can be used to identify the processes occurring in each lake area, leading to a significant improvement in the accuracy of the results. In this sense, a more detailed analysis of the data is required to identify the processes occurring in each lake area. In this regard, we present a long-term spatial-temporal evolution of glacial lakes across the Peruvian Andes and address possible connections with GLOFS events and climate change. We used an open-access satellite-based Landsat 5 image and GTOPO30, and we use a combined using Landsat 5, 30-m GTOPO30

METHODOLOGY
 The pre-processing and image processing procedures for the multi-temporal analysis of glacial lakes were based on an adaptation of the Hydrogeomorphic method (HGM) (Lambert, 2017) and other references (Pielke et al., 2008; Garay et al., 2012; Ryan et al., 2019). The main characteristics of the proposed methodology include:
 - usage of the Landsat collection (30 m spatial resolution, panchromatic image processing temporal detection map), supported by GTOPO30;
 - lake-based processing (using Earth Engine);
 - multi-year validation and field survey.

Figure 2. Methodological flowchart

CONCLUSIONS
 The newly applied methodology based on temporal detection, in the case of lakes, allows the identification of the processes occurring in each lake area and can be used in the 20 Cordilleras of the Peruvian Andes. In general, the results of this study are an important source of information for the analysis of glacial lakes, their evolution, and trends in the last few decades.

AUTHOR INFORMATION **ABSTRACT** **CONTACT AUTHOR** **PRINT** **GET POSTER**

Turpo Cayo E.(4,5); Jara Infantes, H.(1); Melgarejo, E.(1); Tinoco Meyhuay T.(3); Torres, J (1); Caballero, A.(1); Villafane, H.(1); Araujo,J.(1); Yarleque, C.(1); Garay, D. (1); Stephan,H (2); Wood,J (2); Glasser, N.(2); Ryan W.(2); Bennett, J.(2); Shannon, S. (2); Espinoza, R.(5) and De Almeida, C.(6,5)

(1) National Research Institute for Glaciers and Mountain's Ecosystems (INAIGEM); (2) University of Exeter; (3) Santiago Antúnez de Mayolo National University (UNASAM); (4) Institute of the Common Good (IBC); (5) La Molina National Agrarian University, Doctoral Program in Water Resources (PDRH); (6) National Institute for Space Research (INPE)

PRESENTED AT:



ABSTRAC

The effects of climate change are causing a profound reduction in glacier's coverage in the Cordilleras of the Peruvian Andes, giving rise to the development and growth of proglacial lakes. These glacial lakes can represent a significant hazard in the form of Glacier Lake Outburst Floods (GLOF) that have the potential to impact vulnerable populations and infrastructure located downstream of glaciated areas. In this study, we present a long-term spatio-temporal analysis of glacial lakes across the Peruvian Andes and address possible connections with GLOF events and climate change. Accessed and processed using Google Earth Engine and QGIS, this analysis was conducted using Landsat 5, 7 and 8 satellite data (Surface Reflectance Tier 1) acquired between 1984 and 2019. We detected and estimated glacier lake surface area changes using a robust semi-automated method which coupled empirical decision trees, sub-pixel analysis, spectral and topographic indices with many temporal and spatial filters to improve the quality of the glacial lake classifications. Using this classification as a result, we were able to categorize the glacial lakes based on their surface change dynamics over the past three decades. Results highlight a distinct spatial variability in glacial lake changes across the 20 Cordilleras of the Peruvian Andes. Overall, 12 Cordilleras experienced an increase in glacial lake area during the observation period (with increases oscillating from 1% to 38%), whilst the remaining 8 experienced a reduction in area (with decreases ranging from reductions of -2% to -57%). A multi-temporal validation of the semi-automated glacier lake classification method, utilising manually digitized data, revealed glacial lake mapping overall accuracy above >80% across the study area.

METHODOLOGY

The pre-processing and image processing procedures for the multi-temporal mapping of glacial lakes were based on an adaptation of the Mapbiomas project methodology (MapBiomas 2017) and other references (Pekel et al. 2016, Cayo Turpo et al. 2019, Souza et al. 2019). The main characteristics of the proposed methodology include:

- usage of the Landsat collection (30 m spatial resolution),
- pixel-based image processing (empirical decision tree),
- organized by Cordilleras,
- cloud-based computing (Google Earth Engine),
- multi-year validation and field survey.

Figure 1 Methodological flowchart





Figure 2 Empirical decision tree.



RESULTS

According to Figure 4, we showed the evolution of the surface extension of the glacial lakes in the 20 Cordilleras of the Peruvian Andes in the period 1984-2019; where the La Viuda Cordillera has the greatest surface extension. The Cordilleras that presented the greatest increase in the surface of glacial lakes are Ampato, Volcánica, Apolobamba, and Chonta, while the Huaytapallana and Huagoruncho Cordilleras have a decreasing trend. In the case of the Barroso Cordillera, a significant increase was found, related to the activity of damming and regulation of lakes. In the rest of the Cordilleras, there are less representative changes, however, in the lakes individual analysis, important changes were found in most of the studied Cordilleras of the Peruvian Andes.

Figure 3. Classification of glacial lakes based on empirical decision tree.






Figure 4. Glacial lakes area evolution in the 20 Cordilleras of the Peruvian Andes.



The results of image classification of glacial lakes in the period from 1984 to 2019 were subjected to multitemporal validation obtaining a global precision higher than 85%. Besides, it could be to observe that the years with low values in precision are related to problems in the quality of the mosaics and classification thresholds (Figure 5).

Figure 5. Box plot of thematic accuracies reached by Cordilleras in the period from 1984 to 2019.



DISCUSSION

There are several methodologies to classify water bodies in satellites images, of which empirical decision trees are one most used due to their easiness of use and implementation (Souza et al. 2019, MapBiomias Amazonía 2019). However, empirical decision trees can be subjective in the parameters setting at each decision node, leading to temporally inconsistent results, requiring the application of different post-classification filters. For this research, it was decided to use empirical decision trees due to their superior quality when compared to other classifiers such as Random Forest in the Andes mountain range, where there is a general confusion between the classification of water and relief shadows.

The results of the glacial lakes classification in the period from 1984 to 2019 have overall accuracy greater than 85%. This indicates that the methodology is appropriate for the study area. It ought to be noted that the quality of the classification depends on the variables than are used in the decision tree as well as the quality of the annually composed mosaics. Topographic variables like slope and shadow are important variables for the classification of glacial lakes in environments with rugged relief, such as the Andes mountain range. In the results, the Barroso Cordillera shows greater temporal changes, caused by the damming of glacial lakes in view of, the water deficit of this mountain range, which is closely related to the direct and indirect effects of climate change. The results of this study have been compared to the work entitled Global Surface Water (Pekel et al. 2016) and demonstrated to have, greater temporal consistency, which can be explained by the temporal and spatial filters used in this study.

CONCLUSIONS

This study applied a methodology based on empirical decision trees, to the entire Andes mountains. The classification was based on Landsat satellite images and was carried out in the 20 Peruvian Cordilleras in the period 1984-2019. In general, in all Andes mountains, the overall accuracy surpassed 85%. This indicates that the results of this study are an important source of information for the analysis of glacial lakes, their evolution, and trends in the latest three decades.

AUTHOR INFORMATION

Turpo, E¹; Jara, H²; Melgarejo, E²; Tito Tinoco Meyhuay Sr.³; Torres, J²; Caballero, A²; Villafane, H²; Araujo, J²; Yarleque, C²; Garay, D²; Stephan, H⁵; Wood, J⁵; Glasser, N;
Ryan W; Bennett, J; Shannon, S; Espinoza, R and De Almeida, C.

ABSTRACT

The effect of climate change is causing a profound reduction in glacier's coverage in the Cordilleras of the Peruvian Andes, giving rise to the development and growth of proglacial lakes. These glacial lakes can represent a significant hazard in the form of Glacier Lake Outburst Floods (GLOF) that have the potential to impact vulnerable populations and infrastructure located downstream of glaciated areas. In this study, we present a long-term spatio-temporal analysis of glacial lakes across the Peruvian Andes and address possible connections with GLOF events and climate change. Accessed and processed using Google Earth Engine and QGIS, this analysis was conducted using Landsat 5, 7 and 8 satellite data (Surface Reflectance Tier 1) acquired between 1984 to 2019. We detected and estimated glacier lake surface area changes using a robust semi-automated method which coupled empirical decision trees, sub-pixel analysis, spectral and topographic indices with many temporal and spatial filters to improve the quality of the glacial lake classifications. Using this classification as a result, we were able to categorize the glacial lakes based on their surface change dynamics over the past three decades. Results highlight a distinct spatial variability in glacial lake changes across the 20 Cordilleras of the Peruvian Andes. Overall, 12 Cordilleras experienced an increase in glacial lake area during the observation period (ranging from increases of 1% to 38%), whilst the remaining 8 experienced a reduction in area (ranging from reductions of -2% to -57%). A multi-temporal validation of the semi-automated glacier lake classification method presented, utilising manually digitized data, revealed an overall glacial lake mapping accuracy of >80% across the study area.