NASA/ADS

Investigation on 15-days oscillations in the equatorial Spread-F

Show affiliations

Paulino, Igo; Buriti, Ricardo; Paulino, Ana Roberta; Wrasse, Cristiano M.; Medeiros, Amauri; Takahashi, Hisao; Cueva, Ricardo

An intensive study on the start time of equatorial plasma bubble, observed by an all sky imager deployed at São João do Cariri (7.4°S, 36.5°W), and equatorial spread-F, observed by a coherent back scatter deployed at São Luís (2.5°S, 44.2°W) was conducted from 2001 to 2009. Oscillation of 14.5 days was clearly observed in three month (September 2003, October 2005 and January 2008) in the airglow images with amplitudes of 45-60 min. Furthermore, using the data from the radar range time integration (RTI) maps, several events were observed with dominant period of 14.5 days in September 2001, November 2002, January-February 2003, October-December 2005 and November 2008. In such case, the amplitudes of the oscillations were from 3 min up to 60 min. This oscillation could be related to the semidiurnal lunar tide, which appear as an important contributor to the time of occurrence of equatorial spread-F.

Publication:

42nd COSPAR Scientific Assembly. Held 14-22 July 2018, in Pasadena, California, USA, Abstract id. C2.2-31-18.

Pub Date:

July 2018

Bibcode: 2018cosp...42E2606P

Feedback/Corrections? (/feedback/correctabstract?bibcode=2018cosp...42E2606P)